International Journal of Sports, Health and Physical Education

ISSN Print: 2664-7559
ISSN Online: 2664-7567
IJSHPE 2025; 7(2): 09-13
www.physicaleducationjournal.in
Received: 09-05-2025
Accepted: 10-06-2025

Arpita Sharma Pandey
Yoga Coach, SKV, Gwalior, Madhya Pradesh, India

Amit Pandey
Swimming Coach, SDPS, Indore, Madhya Pradesh, India

Dr. Aarti Sharma
Associate Professor, Sage University, Indore, Madhya Pradesh, India

Corresponding Author:
Arpita Sharma Pandey
Yoga Coach, SKV, Gwalior, Madhya Pradesh, India

Comparative study of the effects of breathing techniques in pranayama and swimming on the human body

Arpita Sharma Pandey, Amit Pandey and Aarti Sharma

DOI: https://www.doi.org/10.33545/26647559.2025.v7.i2a.218

Abstract
This research aims to explore and compare the physiological, psychological, and respiratory benefits of breathing techniques used in Pranayama, a yogic practice, and those developed through swimming, a cardiovascular and full-body exercise. Both disciplines involve breath control, yet differ in methodology, intensity, and overall impact on the human body. This comparative study draws on existing scientific literature, observational insights, and biomedical data to highlight the unique and overlapping effects of both practices.

Keywords: Swimming, pranayam, VO_2, RHR, stress level, t-test, sample, PEFR, lung capacity, standard deviation, mean

Introduction
Breathing is a fundamental physiological process essential for life. However, controlled breathing techniques, especially in disciplines like Pranayama and Swimming, serve more than just biological necessity they enhance respiratory function, improve mental health, and build physical endurance. While Pranayama is rooted in ancient Indian yogic traditions, emphasizing slow and mindful breathing, swimming uses rhythmic, controlled breathing to meet the high oxygen demands of vigorous physical activity.
This paper presents a comparative analysis of the effects of these two practices on the human body, highlighting their benefits, differences, and potential applications in health and wellness.

Breathing Techniques: An Overview
Pranayama
Pranayama consists of various controlled breathing techniques such as.
- Anulom-Vilom (alternate nostril breathing).
- Bhramari (humming bee breath).
- Kapalbhati (forced exhalation).
- Kumbhaka (breath retention).

These practices are generally performed in seated, meditative postures with slow, deep inhalations and exhalations through the nose.

Swimming
Swimming requires breath regulation in coordination with physical movement. Breathing is typically rhythmic and involves
- Timed inhalation and exhalation.
- Breath-holding while submerged.
- Rapid exchange of air.

The mouth is often used for inhalation and exhalation due to the face being submerged during strokes.

Physiological effects lung function and respiratory efficiency

- Pranayama increases lung capacity by promoting deep breathing and strengthening the diaphragm. Studies show improvement in Forced Vital Capacity (FVC) and Peak Expiratory Flow Rate (PEFR).
- Swimming also improves lung function significantly due to the body's need to maximize oxygen intake under physical exertion and limited breathing opportunities.

Cardiovascular Health

- Pranayama lowers blood pressure and resting heart rate, enhancing parasympathetic activity.
- Swimming improves cardiovascular endurance and VO_2 max, making the heart more efficient over time.

Psychological Effects

- Pranayama has shown positive effects in reducing anxiety, stress, and symptoms of depression through the regulation of the autonomic nervous system.
- Swimming triggers endorphin release, improving mood and mental focus, and can also act as a meditative activity.

Methodology

The study is based on a comparative literature review and synthesis of peer-reviewed articles, medical journals, and yoga and sports science resources. Physiological and psychological parameters such as lung capacity, heart rate, oxygen consumption (VO_2 max), stress levels, and PEFR were analyzed and compared across both activities.

In the context of our comparative study between Pranayama and swimming, we designed null hypothesis for each physiological parameter.

- **Null Hypothesis (H_0):** There is no significant difference in the mean physiological parameter between individuals practicing Pranayama and those practicing swimming.
- **Alternative Hypothesis (H_1):** There is a significant difference in the mean physiological parameter between the two groups.

Statistical Testing

- We conducted independent samples t-tests for each parameter.
- The p-value tells us whether the data provide enough evidence to reject H_0 in favor of H_1.

To generate the results, the following basic statistical parameters were involved.

Basic parameters used for each group

For each variable (e.g., Lung Capacity, VO_2 Max, etc.) the following were computed.

Parameter	Description
Sample size (n)	Number of participants per group (n = 30)
Mean (μ)	Average value of the parameter for each group
Standard deviation (σ)	Measure of how spread out the data is within each group
Standard error (SE)	Estimate of variability in sample mean: SE = σ / √n
Difference in means	The observed difference between Pranayama and Swimming group means

Degrees of freedom, Estimated using Welch's formula or pooled method for equal variances.

p-value, Probability that observed difference is due to chance, under H_0 |

Following parameters have been considered in study.

- **Lung Capacity (L):** Lung capacity refers to the volume of air your lungs can hold and move in and out during breathing. It's a key indicator of respiratory health, efficiency, and endurance capacity.
- **VO_2 Max (ml/kg/min):** VO_2 Max (maximal oxygen uptake) is one of the most important indicators of cardiovascular fitness. It measures the maximum amount of oxygen our body can utilize per minute during intense exercise.
- **Resting Heart Rate (bpm):** Resting Heart Rate (RHR) is a vital indicator of our cardiovascular fitness and autonomic nervous system balance. It represents the number of times your heart beats per minute when you're at complete rest.
- **Stress Level (scale 1-10):** The stress level (1-10) is a subjective, self-reported measure of an individual's perceived psychological or emotional stress.
- **PEFR (L/min):** Here the PEFR (Peak Expiratory Flow Rate) compares the distribution of PEFR values between the Pranayama and Swimming groups.

Each results are based on: Two sample means, their standard deviations, Sample size and the t-distribution (assuming unequal variances).

Results

T-Test Results: Comparison between Pranayama and Swimming Groups.

Parameter	T-Statistic	P-Value	Significance
Lung Capacity (L)	-5.7208	0.0000	Highly Significant
VO_2 Max (ml/kg/min)	-9.5215	0.0000	Highly Significant
Resting Heart Rate (bpm)	2.8216	0.0065	Significant
Stress Level (1-10 scale)	-3.6404	0.0006	Highly Significant
PEFR (L/min)	-5.1275	0.0000	Highly Significant

The box plots below illustrate the distribution of each parameter for both Pranayama and Swimming groups with Key observations.

- Swimming group consistently shows higher lung capacity, VO_2 max and PEFR, indicating superior respiratory and cardiovascular conditioning.
- Pranayama group demonstrates lower resting heart rate and stress levels, highlighting its effectiveness in relaxation and autonomic regulation.
- All differences were statistically significant ($p<0.01$), confirming meaningful effects from both practices.

For understanding, since the p-value for Resting Heart Rate was 0.0065 < 0.01, we reject H_0 and conclude a significant difference exists. Here is the box plot comparing resting heart rates between the Pranayama and Swimming groups.



Similarly, The PEFR (Peak Expiratory Flow Rate) box plot compares the distribution of PEFR values between the Pranayama and Swimming groups.

The graph generated from our data

Swimming Group

1. **Higher median PEFR:** Indicates stronger lung function due to the aerobic, resistance nature of swimming.
2. **Narrower spread (interquartile range):** Suggests consistency in lung strength across swimmers.

Pranayama Group

1. **Lower median PEFR:** Indicates moderate improvement in lung function.
2. May still show beneficial effects, but less pronounced than swimming.

- **No overlap in interquartile ranges (box edges):** Strong evidence of statistically significant difference between the groups





Outliers (dots beyond whiskers): Show individuals whose PEFR significantly differs from the group trend, which can happen due to individual fitness, technique, or pre-existing conditions.

Similarly





Conclusion

The results demonstrates that swimming leads to a significantly higher PEFR than Pranayama, suggesting that it more effectively enhances lung power and airway capacity. However, Pranayama still improves PEFR, likely through controlled breathing and improved diaphragm use. Significant difference has shown for RHR between groups (P=0.0065), swimmers had lower RHR than pranayama practitioners. This supports the idea that swimming builds cardiovascular efficiency, while pranayama calms the nervous system without major cardiac load.

The 1-10 stress level scale gives a practical insight into participant's emotional states. In our study, it clearly shows that Pranayama has a stronger calming effect on the mind than swimming, which is consistent with yogic science and neurophysiological findings.

In our study, swimming clearly outperformed Pranayama in improving VO_2 Max, making it superior for cardiovascular conditioning. While, swimming enhanced lung capacity more than Pranayama, though both showed benefit swimming due to physical exertion and breath resistance, and Pranayama through controlled, mindful expansion of the lungs.

So In short, Swimming significantly improves cardiovascular and respiratory performance, whereas Pranayama is highly effective in reducing stress and enhancing autonomic regulation. Integrating both practices could offer comprehensive physical and psychological health benefits.

References

1. Jerath R, Edry JW, Barnes VA, Jerath V. Physiology of long pranayamic breathing. Med Hypotheses. 2006;67(3):566-571.
2. Grimby G, Saltin B. The ageing muscle. Clin Physiol. 1983;3(3):209-218.
3. Yadav RK, Das S, Sarkar S, Padhi MM. Effect of yoga on pulmonary function in patients with bronchial asthma. J Altern Complement Med. 2012;18(2):196-201.
4. Mickleborough TD, Lindley MR. The role of nutrition in exercise-induced bronchoconstriction. Curr Sports Med Rep. 2013;12(4):208-217.
5. American College of Sports Medicine. Guidelines for Exercise Testing and Prescription. 10th ed. Philadelphia: Wolters Kluwer, 2021.

6. Saoji AA, Raghavendra BR, Manjunath NK. Effect of yogic breath regulation: A narrative review of scientific evidence. J Ayurveda Integr Med. 2019;10(1):50-58.
7. Pal GK, Velkumary S, Madanmohan. Effect of short-term practice of breathing exercises on autonomic functions in normal human volunteers. Indian J Med Res. 2004;120(2):115-121.
8. Brown RP, Gerbarg PL. Sudarshan Kriya yogic breathing in the treatment of stress, anxiety, and depression. J Altern Complement Med. 2005;11(4):711-717.
9. Tomlin DL, Wenger HA. The relationship between aerobic fitness and recovery from high-intensity intermittent exercise. Sports Med. 2001;31(1):1-11.
10. Zinzen E, Meeusen R. Influence of regular swimming on lung volumes and aerobic capacity in preadolescent girls. Pediatr Exerc Sci. 2001;13(4):361-370.
11. Kjeldsen SE, Helle KB. The importance of physical activity in the prevention of hypertension and cardiovascular disease. Eur. Heart J Suppl. 2014;16(Suppl A):A3-A7.
12. Beaver WL, Wasserman K, Whipp BJ. A new method for detecting the anaerobic threshold by gas exchange. J Appl Physiol. 1986;60(6):2020-2027.
13. Telles S, Desiraju T. Autonomic changes in Brahmakumaris Raja Yoga meditation. Int J Psychophysiol. 1991;11(2):145-151.
14. Dishman RK, Berthoud HR, Booth FW, Cotman CW, Edgerton VR, Fleshner MR, *et al*. Neurobiology of exercise. Obesity (Silver Spring). 2006;14(3):345-356.